

# Tame ordered structures

Antongiulio Fornasiero  
`antongiulio.fornasiero@gmail.com`

University of Münster

Logic Colloquium 2009


# Introduction

- A **definably complete** (DC) structure is a linearly ordered structure  $\mathbb{K}$ , such that every definable subset of  $\mathbb{K}$  has a supremum in  $\mathbb{K} \cup \{\pm\infty\}$ .
- For example, all expansions of  $\mathbb{R}$  and all o-minimal structure are DC.
- All structures in this talk will be assumed to be **DC expansions of ordered fields**.

# Introduction

- A **definably complete** (DC) structure is a linearly ordered structure  $\mathbb{K}$ , such that every definable subset of  $\mathbb{K}$  has a supremum in  $\mathbb{K} \cup \{\pm\infty\}$ .
- For example, all expansions of  $\mathbb{R}$  and all o-minimal structure are DC.
- All structures in this talk will be assumed to be **DC expansions of ordered fields**.
- We do not have yet a precise definition of “tame structures”. We will describe some classes of tame structures (which are DC expansion of fields!) containing the class of o-minimal structures.
- Tame non-DC ordered structure (e.g., weakly o-minimal structures) are outside the scope of this talk.


# The overall picture



# Contents

- 1 Locally o-minimal structures
- 2 D-minimal structures
- 3 Structures satisfying Uniform Finiteness
- 4 Structures satisfying Pseudo-Finiteness
- 5 Baire structures
- 6 Dimension in d-minimal structures

# The overall picture



# Locally o-minimal structures

## Definition

$\mathbb{K}$  is **locally o-minimal** if it is o-minimal “around every point”.

That is, for every  $X$  definable subset of  $\mathbb{K}$  and every  $a \in \mathbb{K}$ , there exists  $0 < \varepsilon \in \mathbb{K}$ , such that  $(a, a + \varepsilon)$  is either disjoint from  $X$ , or contained in  $X$ .

## Example

Ultraproducts of o-minimal structures.

## Remark

If  $\mathbb{K}$  expands  $\mathbb{R}$ , then  $\mathbb{K}$  is locally o-minimal iff it is o-minimal.

The same remains true if we only ask that  $\mathbb{K}$  is Archimedean.

# Properties of locally o-minimal structures

Many facts from the theory of o-minimal structures remain true for locally o-minimal ones.

**Monotonicity theorem** Given a definable function  $f : \mathbb{K} \rightarrow \mathbb{K}$ , there exists a **pseudo-finite** set  $I$  such that, on each intervals of  $K \setminus I$ ,  $f$  is continuous and either constant or strictly monotone. Same with  $C^p$  instead of continuous.

**Constructibility** Every definable subset of  $\mathbb{K}^n$  is **constructible** (finite Boolean combination of definable open sets).

**Definable Choice**  $\mathbb{K}$  has definable Skolem functions and elimination of imaginaries

**Growth dichotomy**  $\mathbb{K}$  is either power-bounded, or defines an exponential function.



# Properties no longer true for locally o-minimal structures

**Exchange principle** The algebraic closure does not satisfy the exchange principle for the algebraic closure.


## Theorem (A. Dolich)

*If a locally o-minimal theory  $T$  satisfies the exchange principle, then  $T$  is o-minimal.*

**Rosiness** A locally o-minimal non o-minimal theory is not rosy.

**NIP** There exists an ultraproduct of o-minimal structures (and hence locally o-minimal) which satisfies the Independence Property.

# The overall picture



# D-minimal structures

## Definition (C. Miller)

$\mathbb{K}$  is **d-minimal** if, for every  $\mathbb{K}'$  elementary extension of  $\mathbb{K}$ , and every  $X$  definable subset of  $\mathbb{K}'$ ,  $X$  is the union of an open set and finitely many **discrete** sets.

## Example (van den Dries)

$(\mathbb{R}, +, \cdot, <, 2^{\mathbb{Z}})$  is d-minimal, where  $2^{\mathbb{Z}}$  is a predicate denoting the set of real integer powers of 2.

## Properties of d-minimal structures

Many of the properties of locally minimal structures remain true for d-minimal ones. Let  $\mathbb{K}$  be a d-minimal structure.


**Monotonicity theorem** Given a definable function  $f : \mathbb{K} \rightarrow \mathbb{K}$ , there exists a definable set  $I$  which is **closed** and with **empty interior**, and such that, on each intervals of  $K \setminus I$ ,  $f$  is continuous and either constant or strictly monotone. Same with  $C^p$  instead of continuous.

**Constructibility** Every definable subset of  $\mathbb{K}^n$  is **constructible**.

**Definable Choice**  $\mathbb{K}$  has definable Skolem functions and elimination of imaginaries

**(Un)rosiness** If  $\mathbb{K}$  is not o-minimal, then it is not rosy.

# The overall picture



# Open core and dense pairs

## Definition

The **open core** of  $\mathbb{K}$  is the reduct of  $\mathbb{K}$  generated by all open definable subsets of  $\mathbb{K}$ .

## Definition

Given a theory  $T$  (expanding RCF), let  $T^d$  is the theory **dense pairs** of models of  $T$ : that is, the theory of pairs  $(A, B)$ , such that

- 1  $A < B \models T$ ;
- 2  $A \neq B$ ;
- 3  $A$  is dense in  $B$ :  
 $\forall b < b' \in B \exists a \in A (b < a < b')$ .

# Structure satisfying Uniform Finiteness

## Definition

$\mathbb{K}$  satisfies **Uniform Finiteness (UF)** if it eliminates the quantifier “there exist infinitely many”.

## Theorem (Dolich, Miller, Steinhorn)

$K$  satisfies UF iff its open core is o-minimal.


## Theorem (Dries, DMS)

If  $T$  is an o-minimal theory, then  $T^d$  is complete, it satisfies DC and UF, and the open core of  $T^d$  is  $T$ .

## Theorem (F.)

If  $T$  is a d-minimal theory, then  $T^d$  is complete. The Cauchy completion  $\tilde{\mathbb{K}}$  of a d-minimal structure  $\mathbb{K}$  has a unique structure such that  $\mathbb{K} \leq \tilde{\mathbb{K}}$ ; therefore,  $T^d$  is also consistent.

# The overall picture





# Pseudo-finite sets

## Definition

$X \subseteq \mathbb{K}$  is **pseudo-finite** if it is definable, discrete, closed, and bounded.

## Remark

If  $\mathbb{K}$  is an expansion of the real line, then any pseudo-finite subset of  $\mathbb{K}^n$  is finite.

## Lemma

*If  $X \subset \mathbb{K}^n$  is pseudo-finite and  $f : \mathbb{K}^n \rightarrow \mathbb{K}^m$  is definable, then  $f(X)$  is also pseudo-finite.*

## Pseudo-finite sets, continued

### Example

If  ${}^*\mathbb{R}$  is a non-standard model of analysis, and  $X \subset {}^*\mathbb{R}$  is definable, then  $X$  is pseudo-finite iff its cardinality is a (standard or not) natural number.

### Remark

$\mathbb{K}$  satisfies UF iff every pseudo-finite subset is finite.

# Structures satisfying Pseudo-finiteness

## Lemma

$\mathbb{K}$  satisfies **Uniform Finiteness (UF)** iff every definable discrete subset of  $\mathbb{K}$  is finite.

# Structures satisfying Pseudo-finiteness

## Definition

$\mathbb{K}$  satisfies **Pseudo-Finiteness** ( $\Psi F$ ) iff every definable discrete subset of  $\mathbb{K}$  is pseudo-finite (that is, closed and bounded).

## Lemma

- 1 If  $\mathbb{K}$  satisfies UF, then it satisfies  $\Psi F$ .
- 2  $\Psi F$  is an elementary property: in particular, ultraproducts of structures with  $\Psi F$  do satisfy  $\Psi F$ .

# Structures satisfying Pseudo-finiteness

## Definition

$\mathbb{K}$  satisfies **Pseudo-Finiteness** ( $\Psi F$ ) iff every definable discrete subset of  $\mathbb{K}$  is pseudo-finite (that is, closed and bounded).

## Lemma

- 1 *If  $\mathbb{K}$  satisfies UF, then it satisfies  $\Psi F$ .*
- 2  *$\Psi F$  is an elementary property: in particular, ultraproducts of structures with  $\Psi F$  do satisfy  $\Psi F$ .*

## Theorem (DMS)

$\mathbb{K}$  satisfies UF iff its open core is o-minimal.

# Structures satisfying Pseudo-finiteness

## Definition

$\mathbb{K}$  satisfies **Pseudo-Finiteness** ( $\Psi F$ ) iff every definable discrete subset of  $\mathbb{K}$  is pseudo-finite (that is, closed and bounded).


## Lemma

- 1 *If  $\mathbb{K}$  satisfies UF, then it satisfies  $\Psi F$ .*
- 2  *$\Psi F$  is an elementary property: in particular, ultraproducts of structures with  $\Psi F$  do satisfy  $\Psi F$ .*


## Theorem (F.)

$\mathbb{K}$  satisfies  $\Psi F$  iff its open core is locally o-minimal.

# The overall picture




# The overall picture







## The overall picture




# The overall picture




# The overall picture



## The overall picture



# The overall picture



# Baire structures

## Definition

A subset  $X$  of  $\mathbb{K}^n$  is **nowhere dense** if the closure of  $X$  has empty interior:  
 $\overset{\circ}{\overline{X}} = \emptyset$ .

$\mathbb{K}$  is definably **Baire** if it is not the union of an increasing definable family of nowhere dense subsets.

## Definition

$X \subseteq \mathbb{K}^n$  is **definably meager** if  $X$  is the union of an increasing definable family of nowhere dense subsets.

## Remark

$\mathbb{K}$  is (definably) Baire iff it is not definably meager.

Being Baire is a first-order property, and hence preserved under elementary equivalence and ultraproducts.

## Baire structures, continued

### Examples

- Every expansion of  $\mathbb{R}$  is Baire.
- Every d-minimal structure is Baire.
- Every structure satisfying  $\Psi F$  is Baire.

### Lemma

If  $\mathbb{K}$  defines a discrete subring  $N$ , then  $\mathbb{K}$  is Baire.

### Proof.

$N$  is a model of first-order Peano Axioms; hence, one can simulate in  $N$  a proof of the “classical” Baire Category Theorem.  $\square$

### Conjecture

Every DC expansion of an ordered field is Baire.

# Dimension

## Definition

Let  $X \subseteq \mathbb{K}^n$ . The **dimension** of  $X$  is  $\dim(X)$ , the largest integer  $d$  such that there exists a coordinate space  $L$  of dimension  $d$ , such that  $\Pi_L(X)$  has non-empty interior, where  $\Pi_L$  is the projection from  $\mathbb{K}^n$  to  $L$ .

## Remark

- $\dim$  is monotone;
- if  $X \subseteq \mathbb{K}^n$ , then  $\dim(X) \leq n$ .

It is not true in general that  $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$ .



# Dimension in d-minimal structures

Let  $\mathbb{K}$  be **d-minimal**, and  $X$  and  $Y$  be definable sets.

## Lemma (1)

- $\dim(X \cup Y) = \max(\dim(X), \dim(Y))$ ;
- if  $X \subseteq \mathbb{K}^{n+m}$ ,  $Y := \Pi_n^{n+m}(X)$ ,  $\dim(Y) = d$ , and  $\dim(X_a) = k$  for every  $a \in Y$ , then  $\dim(X) = d + k$ .

## Lemma

- $\dim(\overline{X}) = \dim X$ ;
- *Sard's Lemma*: if  $f : \mathbb{K}^n \rightarrow \mathbb{K}^m$  is  $C^1$  function definable in  $\mathbb{K}$ , then the set of **critical points** of  $f$  has empty interior.
- If moreover  $\mathbb{K}$  is **locally o-minimal**, then  $\dim(\partial X) < \dim(X)$ .

# Dimension functions

Other kind of structures have well-behaved notions of dimension too.

## Definition

A **dimension function** (on some structure  $\mathbb{K}$ ) is a function  $\dim$  from definable sets into natural numbers, such that

- 1  $\dim$  is monotone;
- 2 the dimension of any singleton is 0;
- 3  $\dim(\mathbb{K}) = 1$ ;
- 4  $\dim$  is additive;
- 5 for every integer  $d$  and definable set  $X \subseteq \mathbb{K}^{n+m}$ ,
  - a. the set  $\{a \in \mathbb{K}^n : \dim(X_a) = d\}$  is definable;
  - b. let  $Y := \Pi_n^{n+m}(X)$ ; if  $\dim(Y) = d$  and  $\dim(X_a) = k$  for every  $a \in Y$ , then  $\dim(X) = d + k$ .