

Diamond, non-saturation, and weak square principles

*Logic Colloquium 2009
(European ASL meeting)*

02-Aug-09, Sofia

Assaf Rinot
Tel-Aviv University

<http://www.tau.ac.il/~rinot>

Diamond on successor cardinals

Definition (Jensen, '72). For an infinite cardinal, λ , and a stationary set $S \subseteq \lambda^+$, $\diamond(S)$ asserts the existence of a sequence $\langle A_\alpha \mid \alpha \in S \rangle$ such that $\{\alpha \in S \mid A \cap \alpha = A_\alpha\}$ is stationary for all $A \subseteq \lambda^+$.

Fact. For $S \subseteq \lambda^+$, $\diamond_S \Rightarrow \diamond_{\lambda^+} \Rightarrow 2^\lambda = \lambda^+$.

Question. Given a stationary, $S \subseteq \lambda^+$,
Does $2^\lambda = \lambda^+ \Rightarrow \diamond(S)$?

A related concept

Fact. $\diamond(S)$ entails that $\text{NS}_{\lambda^+} \upharpoonright S$ is *non-saturated*.

That is, there exists a family of λ^{++} many stationary subsets of S , whose pairwise intersection is nonstationary.

Proof. Let $\langle A_\alpha \mid \alpha \in S \rangle$ witness $\diamond(S)$. Denote $S_A = \{\alpha \mid A \cap \alpha = A_\alpha\}$. Then $\{S_A \mid A \subseteq \lambda^+\}$ exemplifies the non-saturation of $\text{NS}_{\lambda^+} \upharpoonright S$. ■

Question. Given a stationary, $S \subseteq \lambda^+$,
Must $\text{NS}_{\lambda^+} \upharpoonright S$ be non-saturated?

Two negative results. $\lambda = \omega$

Theorem (Jensen, '74). It is consistent that CH holds, while $\diamond(\omega_1)$ fails.

Theorem (Steel-Van Wesep, '82). Suppose that V is a model of “ZF + $AD_{\mathbb{R}}$ + Θ is regular”.

Then, there is a forcing extension which is a model of ZFC, in which NS_{ω_1} is saturated.

Remark. By Later work of Shelah and Jensen-Steel, the saturation of NS_{ω_1} is equiconsistent with the existence of a single Woodin cardinal.

Two positive results. $\lambda > \omega$

Denote $E_{\neq \kappa}^{\lambda^+} := \{\delta < \lambda^+ \mid \text{cf}(\delta) \neq \kappa\}$.

Theorem (Shelah, '90s). If λ is an uncountable cardinal, and S is a stationary subset of $E_{\neq \text{cf}(\lambda)}^{\lambda^+}$, then $\text{NS}_{\lambda^+} \upharpoonright S$ is non-saturated.

A continuous effort of 30 years recently culminated in:

Theorem (Shelah, 2007). If λ is an uncountable cardinal, and S is a stationary subset of $E_{\neq \text{cf}(\lambda)}^{\lambda^+}$, then $2^\lambda = \lambda^+ \Rightarrow \diamond(S)$.

The critical cofinality. $\lambda = \text{cf}(\lambda)$

Denote $E_{\kappa}^{\lambda^+} := \{\delta < \lambda^+ \mid \text{cf}(\delta) = \kappa\}$.

Theorem (Shelah, '80). For every regular uncountable cardinal, λ , it is consistent that:

$$\text{GCH} + \neg \diamond(E_{\lambda}^{\lambda^+}) .$$

Theorem (Woodin, '80s). For every regular uncountable cardinal, λ , having a huge cardinal above it, in some $< \lambda$ -closed forcing extension:

$$\text{NS}_{\lambda^+} \upharpoonright S \text{ saturated, for some stationary } S \subseteq E_{\lambda}^{\lambda^+} .$$

The critical cofinality. $\lambda > \text{cf}(\lambda)$

Def. $S \subseteq \lambda^+$ *reflects* iff the following set is stationary:

$$\text{Tr}(S) := \{\gamma < \lambda^+ \mid \text{cf}(\gamma) > \omega, S \cap \gamma \text{ is stationary}\}.$$

Theorem (Shelah, '84). For every singular cardinal, λ , in some cofinality-preserving forcing extension:

$\text{GCH}^+ \rightarrow \diamond(S)$ for some **non-reflecting** stationary set $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$.

Theorem (Foreman, '83). For every singular cardinal, λ , having a supercompact cardinal above it, and an almost-huge cardinal above that supercompact, in some λ -preserving forcing extension:

$\text{NS}_{\lambda^+} \upharpoonright S$ saturated, for a **non-reflecting** stationary $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$.

Questions

Question 1. Suppose λ is a singular cardinal.

Must $2^\lambda = \lambda^+ \Rightarrow \diamond_S$ for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects?

Question 2. Suppose λ is a singular cardinal.

Must $\text{NS}_{\lambda^+} \upharpoonright S$ be non-saturated for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects?

Question 3. Can $\text{NS}_{\omega_2} \upharpoonright E_{\omega_1}^{\omega_2}$ be saturated?

Some answers



Diamond and reflecting sets

A partial **affirmative** answer to Question 1 is provided by Shelah and Zeman, as follows.

Theorem (Shelah, '84). If $2^\lambda = \lambda^+$ for a **strong limit** singular cardinal λ , and \square_λ^* holds, then $\diamond(S)$ for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects.

Theorem (Zeman, 2008). If $2^\lambda = \lambda^+$ for a singular cardinal λ , and \square_λ^* holds, then $\diamond(S)$ for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects.

Weak Square

Definition (Jensen '72). \square_λ^* asserts the existence of a sequence $\langle \mathcal{P}_\alpha \mid \alpha < \lambda^+ \rangle$ such that:

1. $\mathcal{P}_\alpha \subseteq [\alpha]^{<\lambda}$ and $|\mathcal{P}_\alpha| = \lambda$ for all $\alpha < \lambda^+$;
2. for every limit $\gamma < \lambda^+$, there exists
a club $C_\gamma \subseteq \gamma$ satisfying:

$$C_\gamma \cap \alpha \in \mathcal{P}_\alpha \text{ for all } \alpha < \gamma.$$

The approachability ideal

Definition (Shelah). A set T is in $I[\lambda^+]$ iff:

1. $T \subseteq \lambda^+$;
2. there exists a sequence $\langle \mathcal{P}_\alpha \mid \alpha < \lambda^+ \rangle$ such that:
 - 2.1. $\mathcal{P}_\alpha \subseteq [\alpha]^{<\lambda}$ and $|\mathcal{P}_\alpha| = \lambda$ for all $\alpha < \lambda^+$;
 - 2.2. for almost all $\gamma \in T$, there exists an unbounded $A_\gamma \subseteq \gamma$ satisfying:

$$A_\gamma \cap \alpha \in \bigcup_{\beta < \gamma} \mathcal{P}_\beta \text{ for all } \gamma < \alpha.$$

A relative of approachability ideal

Definition. Given $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$, a set T is in $I[S; \lambda]$ iff:

1. $T \subseteq \text{Tr}(S)$;
2. there exists a sequence $\langle \mathcal{P}_\alpha \mid \alpha < \lambda^+ \rangle$ such that:
 - 2.1. $\mathcal{P}_\alpha \subseteq [\alpha]^{<\lambda}$ and $|\mathcal{P}_\alpha| = \lambda$ for all $\alpha < \lambda^+$;
 - 2.2. for almost all $\gamma \in T$, there exists a stationary $S_\gamma \subseteq S \cap \gamma$ satisfying:
$$S_\gamma \cap \alpha \in \bigcup \{ \mathcal{P}(X) \mid X \in \mathcal{P}_\alpha \} \text{ for all } \alpha < \gamma$$

Remark. If λ is SSL, then $I[S; \lambda] \subseteq I[\lambda^+]$.

A comparison with weak square

Let λ denote a singular cardinal, and let $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$.

Observation. If $I[S; \lambda]$ contains a stationary set, then S reflects.

Proposition. Assume \square_{λ}^* . If S reflects, then $I[S; \lambda]$ contains a stationary set.

Theorem. It is relatively consistent with the existence of a supercompact cardinal that \square_{λ}^* fails, while $I[S; \lambda]$ contains a stationary set for every stationary $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$.

Answering question 1

Improving the Shelah-Zeman theorem, we have:

Theorem. Suppose λ is a singular cardinal, $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$;
If $I[S; \lambda]$ contains a stat. set, then $2^\lambda = \lambda^+ \Rightarrow \diamond(S)$.

Answering Question 1 **in the negative**, while establishing that the above improvement is optimal, we have:

Theorem (Gitik-R.). It is relatively consistent with the existence of a supercompact cardinal that:

- (1) GCH holds;
- (2) $\aleph_{\omega+1} \in I[\aleph_{\omega+1}]$;
- (3) Every stationary subset of $E_\omega^{\aleph_{\omega+1}}$ reflects;
- (4) $\diamond(S)$ fails, for some (reflecting) $S \subseteq E_\omega^{\aleph_{\omega+1}}$.

Stationary Approachability Property

Theorem (Shelah, '84). If $2^\lambda = \lambda^+$ for a strong limit singular cardinal λ , and \square_λ^* holds, then $\diamond(S)$ for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects.

Theorem. If $2^\lambda = \lambda^+$ for a strong limit singular cardinal λ , and \square_λ^* holds, **and every stationary subset of $E_{\text{cf}(\lambda)}^{\lambda^+}$ reflects**, then, moreover, $\diamond^*(\lambda^+)$ holds.

Definition. SAP_λ denote the assertion that $I[S; \lambda]$ contains a stationary set for every $S \subseteq E_{\text{cf}(\lambda)}^{\lambda^+}$ that reflects.

Theorem. Replacing \square_λ^* with SAP_λ is impossible, in the sense that the conclusion would fail to hold. (obtained by forcing over a model with a supercompact.)

Summary: Square vs. Diamond

Let Refl_λ denote the assertion that every stationary subset of $E_{\text{cf}(\lambda)}^{\lambda^+}$ reflects.

Then, for λ singular, we have:

1. $\text{GCH} + \square_\lambda^* \not\Rightarrow \diamond^*(\lambda^+)$;
2. $\text{GCH} + \text{Refl}_\lambda + \square_\lambda^* \Rightarrow \diamond^*(\lambda^+)$;
3. $\text{GCH} + \text{Refl}_\lambda + \text{SAP}_\lambda \not\Rightarrow \diamond^*(\lambda^+)$;
4. $\text{GCH} + \text{Refl}_\lambda + \text{SAP}_\lambda \Rightarrow \diamond(S)$ for every stat. $S \subseteq \lambda^+$;
5. $\text{GCH} + \text{Refl}_\lambda + \text{AP}_\lambda \not\Rightarrow \diamond(S)$ for every stat. $S \subseteq \lambda^+$.

Remark. AP_λ asserts that $\lambda^+ \in I[\lambda^+]$.

Around question 2

Let λ denote a singular cardinal, and $S \subseteq E_{\text{cf}(\lambda)}^{\lambda+}$.

Theorem (Gitik-Shelah, '97).

$\text{NS}_{\lambda+} \upharpoonright E_{\text{cf}(\lambda)}^{\lambda+}$ is non-saturated.

Theorem (Krueger, 2003).

If $\text{NS}_{\lambda+} \upharpoonright S$ is saturated, then S is co-fat.

Theorem. If $\text{NS}_{\lambda+} \upharpoonright S$ is saturated, then $I[S; \lambda]$ does not contain a stationary set.

In particular, SAP_{λ} (and hence \square_{λ}^*) imposes a positive answer to Question 2.

The effect of smaller cardinals



A shift in focus

Instead of studying the validity of $\diamond(S)$ (or saturation), we now focus on finding sufficient conditions for $I[S; \lambda]$ to contain a stationary set.

This yields a linkage between virtually unrelated objects.

Theorem. Assume GCH and that κ is an uncountable cardinal with no κ^+ -Souslin trees.

Then $\diamond(E_{\text{cf}(\lambda)}^{\lambda^+})$ holds for the class of singular cardinals λ of cofinality κ .

let us explain how small cardinals effects $\lambda..$

The effect of smaller cardinals, I

Definition. Assume $\theta > \kappa > \omega$ are regular cardinals.

$R_1(\theta, \kappa)$ asserts that for every function $f : E_{<\kappa}^\theta \rightarrow \kappa$, there exists some $j < \kappa$ such that:

$\{\delta \in E_\kappa^\theta \mid f^{-1}[j] \cap \delta \text{ is stationary}\}$ is stationary.

Facts. 1. $\square_\kappa \Rightarrow \neg R_1(\kappa^+, \kappa)$;

2. every stationary subset of $E_\kappa^{\kappa^{++}}$ reflects $\Rightarrow R_1(\kappa^{++}, \kappa^+)$;

3. By Harrington-Shelah '85, $R_1(\aleph_2, \aleph_1)$ is equiconsistent with the existence of a Mahlo cardinal.

The effect of smaller cardinals, II

Theorem. Suppose $\lambda > \text{cf}(\lambda) = \kappa > \omega$;
If there exists a regular $\theta \in (\kappa, \lambda)$ such that $R_1(\theta, \kappa)$
holds, then $I[E_{\text{cf}(\lambda)}^{\lambda^+}; \lambda]$ contains a stationary set.

Corollary. Suppose κ is a regular cardinal and every
stationary subset of $E_{\kappa}^{\kappa^{++}}$ reflects.

Then $2^\lambda = \lambda^+ \Rightarrow \diamond(E_{\text{cf}(\lambda)}^{\lambda^+})$ for the class of singular
cardinals λ of cofinality κ^+ .

Corollary. Assume PFA^+ ;
 $\diamond(E_{\text{cf}(\lambda)}^{\lambda^+})$ holds for every λ strong limit of cofinality ω_1 .

The effect of smaller cardinals, III

Definition. Assume $\theta > \kappa > \omega$ are regular cardinals.

$R_2(\theta, \kappa)$ asserts that for every function $f : E_{<\kappa}^\theta \rightarrow \kappa$, there exists some $j < \kappa$ such that:

$\{\delta \in E_\kappa^\theta \mid f^{-1}[j] \cap \delta \text{ is non-stationary}\}$ is non-stationary.

Facts. 1. $R_2(\theta, \kappa) \Rightarrow R_1(\theta, \kappa)$ and hence the strength of $R_2(\kappa^+, \kappa)$ is at least of a Mahlo cardinal.

2. By Magidor '82, $R_2(\aleph_2, \aleph_1)$ is relatively consistent with the existence of a weakly compact cardinal.

Remark. The exact strength of $R_2(\aleph_2, \aleph_1)$ is unknown.

The effect of smaller cardinals, IV

Theorem. Suppose $\lambda > \text{cf}(\lambda) = \kappa > \omega$;
If there exists a regular $\theta \in (\kappa, \lambda)$ such that $R_2(\theta, \kappa)$
holds, then $\text{Tr}(S) \cap E_\theta^{\lambda^+} \in I[S; \lambda]$ for every $S \subseteq \lambda^+$.

Corollary. Suppose $R_2(\theta, \kappa)$ holds.

For every sing. cardinal λ of cofinality κ with $2^\lambda = \lambda^+$:

$\diamond(S)$ holds whenever $\text{Tr}(S) \cap E_\theta^{\lambda^+}$ is stationary.

Remark. The $R_2(\cdot, \cdot)$ proof resembles the one of an analogous theorem by Viale-Sharon concerning the weak approachability ideal. The $R_1(\cdot, \cdot)$ proof builds on a fundamental fact from Shelah's *pcf* theory.

Generalized stationary sets



The sup function, I

Definition. A set $\mathcal{X} \subseteq \mathcal{P}(\lambda^+)$ is *stationary* (in the generalized sense) iff for every $f : [\lambda^+]^{<\omega} \rightarrow \lambda^+$, there exists some $X \in \mathcal{X}$ such that $f : [X]^{<\omega} \subseteq X$.

Question (König-Larson-Yoshinobu). Let λ denote an infinite cardinal. Is it possible to prove in ZFC that every stationary $\mathcal{B} \subseteq [\lambda^+]^\omega$ can be thinned out to a stationary $\mathcal{A} \subseteq \mathcal{B}$ on which the sup-function is injective?

The sup function, II

Question (König-Larson-Yoshinobu). Let λ denote an infinite cardinal. Is it possible to prove in ZFC that every stationary $\mathcal{B} \subseteq [\lambda^+]^\omega$ can be thinned out to a stationary $\mathcal{A} \subseteq \mathcal{B}$ on which the sup-function is injective?

Proposition. If $\mathcal{A} \subseteq [\lambda^+]^\omega$ is a stationary set on which the sup-function is injective, then $\text{cf}([\lambda^+]^\omega, \subseteq) = \lambda^+$.

In particular, if the SCH fails, then we get a counterexample to the above question. But what can one say in the context of GCH?

► It turns out that diamond helps..

The sup function, III

Theorem. Suppose λ is a cardinal, $2^\lambda = \lambda^+$.

For a stationary $S \subseteq E_{<\lambda}^{\lambda^+}$, TFAE:

- 1) $\diamond(S)$;
- 2) there exists a stationary $\mathcal{X} \subseteq [\lambda^+]^{<\lambda}$, with $\{\sup(X) \mid X \in \mathcal{X}\} \subseteq S$, on which the sup-function is injective.

Corollary. A negative answer to the K-L-Y question.

Proof. Work in a model of GCH and there exists $S \subseteq E_\omega^{\aleph_{\omega+1}}$ on which $\diamond(S)$ fails.

Put $\mathcal{B} := \{X \in [\aleph_{\omega+1}]^\omega \mid \sup(X) \in S\}$. Then \mathcal{B} is stationary, and no stationary subset of \mathcal{B} can be refined to a set on which the sup-function is injective. ■

A related result

Theorem. Let λ denote an infinite cardinal.

Suppose $\mathcal{X} \subseteq [\lambda^+]^{<\lambda}$ is a stationary set on which the sup-function is $(\leq \lambda)$ -to-1. Put $S := \{\text{sup}(X) \mid X \in \mathcal{X}\}$. Then $\text{NS}_{\lambda^+} \upharpoonright S$ is non-saturated.

λ^+ -guessing



A very weak consequence of $\diamond(E_{\text{cf}(\lambda)}^{\lambda^+})$

Definition. For a function $f : \lambda^+ \rightarrow \text{cf}(\lambda)$, let κ_f denote the minimal cardinality of a family $\mathcal{P} \subseteq [\lambda^+]^{\text{cf}(\lambda)}$ with the following property.

For all $Z \subseteq \lambda^+$ such that $\bigwedge_{\beta < \text{cf}(\lambda)} |Z \cap f^{-1}\{\beta\}| = \lambda^+$, there exist some $a \in \mathcal{P}$ with $\sup(f[a \cap Z]) = \text{cf}(\lambda)$.*

Definition. For a singular cardinal λ , we say that λ^+ -guessing holds iff $\kappa_f \leq \lambda^+$ for all $f \in {}^{\lambda^+}\text{cf}(\lambda)$.

*Note that if λ is SSL, then we may assume that \mathcal{P} is closed under taking subsets. Thus, we may moreover demand the existence of $a \in \mathcal{P}$ such that $a \subseteq Z$ and $f \upharpoonright a$ is injective.

the failure of λ^+ -guessing

Theorem (Džamonja-Shelah, 2000). It is relatively consistent with the existence of a supercompact cardinal that there exist a strong limit singular cardinal, λ , and a function $f : \lambda^+ \rightarrow \text{cf}(\lambda)$ such that $\kappa_f = 2^\lambda > \lambda^+$.

Theorem. Suppose λ is a strong limit singular; then:

$$\{\kappa_f \mid f \in {}^{\lambda^+}\text{cf}(\lambda)\} = \{0, 2^\lambda\}.$$

Corollary. For a strong limit singular cardinal, λ , TFAE:

- 1) λ^+ -guessing;
- 2) $\diamond^*(E_{\neq \text{cf}(\lambda)}^{\lambda^+})$.

a fundamental cardinal arithmetic statement in disguise

Theorem. The following are equivalent:

1. λ^+ -guessing holds for every singular cardinal, λ ;
2. Shelah's Strong Hypothesis, i.e.,
$$\text{cf}([\lambda]^{\text{cf}(\lambda)}, \subseteq) = \lambda^+$$
 for every singular cardinal, λ .
3. Every first-countable topol. space whose density is a regular cardinal, κ , enjoys the following reflection: if every **separable subspace** is of size $\leq \kappa$, then the whole space is of size $\leq \kappa$.

Open problems



Open problems

Let λ denote a singular cardinal.

Question I. Does $2^\lambda = \lambda^+$ entail $\diamond(E_{\text{cf}(\lambda)}^{\lambda^+})$?

Question II. Must there exist a stationary subset of $E_{>\text{cf}(\lambda)}^{\lambda^+}$ that carries a partial (weak) square sequence?

Question III. Is “ NS_{ω_1} saturated” consistent with CH?

Thank you!

